



Review

Is there a path beyond BOLD? Molecular imaging of brain function

Alan P. Koretsky*

Laboratory of Functional and Molecular Imaging, National Institutes of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD 20892, USA

ARTICLE INFO

Article history:

Accepted 27 February 2012

Available online 3 March 2012

Keywords:

Manganese enhanced MRI

MRI calcium indicators

MRI gene expression reporters

MRI detection of neurotransmitters

MRI detection of membrane potential

ABSTRACT

The dependence of BOLD on neuro-vascular coupling leaves it many biological steps removed from direct monitoring of neural function. MRI based approaches have been developed aimed at reporting more directly on brain function. These include: manganese enhanced MRI as a surrogate for calcium ion influx; agents responsive to calcium concentrations; approaches to measure membrane potential; agents to measure neurotransmitters; and strategies to measure gene expression. This work has led to clever design of molecular imaging tools and many contributions to studies of brain function in animal models. However, a robust approach that has potential to get MRI closer to neurons in the human brain has not yet emerged.

© 2012 Published by Elsevier Inc.

Contents

Developing MR for measuring brain function prior to BOLD	1208
Direct measurement of membrane potential: the holy grail	1209
Calcium dynamics using MRI.	1209
Measurement of intracellular calcium	1209
Manganese Enhanced MRI (MEMRI) as a surrogate for calcium influx	1210
MRI agents to measure neurotransmitter release	1211
MRI reporters of gene expression.	1211
Conclusion	1212
Acknowledgments	1213
References	1213

Developing MR for measuring brain function prior to BOLD

It can be argued that 1990 was a landmark year for MRI of the brain. For this NeuroImage issue on the 20th anniversary of fMRI, Seiji Ogawa's 1990 publications describing BOLD are most important (Ogawa et al., 1990a, Ogawa and Lee, 1990, Ogawa et al., 1990b). That same year Moseley and colleagues published the observation that the apparent diffusion coefficient of water decreased during stroke (Moseley et al., 1990a) and that the apparent diffusion coefficient of water is anisotropic in white matter (Moseley et al., 1990b), the basis of diffusion tensor imaging. However, throughout the 1980s there was growing interest in measuring parameters with magnetic resonance that had relevance to brain function. This

included the pioneering spectroscopy studies on brain from the Chance (Schnall et al., 1987), Radda (Radda, 1992), and Shulman (Pritchard and Shulman, 1986) laboratories. This work primarily focused on phosphorus energy metabolism, ^{13}C measurements of glucose metabolism, and ^1H studies of metabolites. In 1983, Smith et al. published a paper describing a ^{19}F labeled calcium chelator for MR measurements of calcium concentration (Smith et al., 1983) that was used to monitor function in brain slices (Bador-Goffer et al., 1990). This agent was inspired by the early demonstration of fluorescent based calcium indicators (Tsien, 1980). Fluorescent calcium indicators have grown into an important tool for measuring neural function. The recent success of genetically encoded, fluorescent calcium indicators will help sustain increasingly sophisticated studies of neural function (Looger and Griesbeck, 2012; Miyawaki et al., 1997). Prior to 1990, it was also being recognized that ^{19}F perfluorocarbon emulsions could be used to measure oxygen with MRI in vivo (Clark et al., 1984). Ackerman and colleagues explicitly set out to

* LFMI, NINDS, Building 10, Rm B1D728, 10 Center Drive, Bethesda, MD 20892, USA. Fax: +1 301 480 2558.

E-mail address: koretskya@ninds.nih.gov.

measure tissue function in analogy to PET measurements of deoxyglucose and regional blood flow (Ackerman et al., 1987; Deuel et al., 1985). Interest had also been developing to use oxygen-17 to measure blood flow and oxygen consumption (Arai et al., 1990).

In 1986, the discovery that small iron oxide particles were potent MRI contrast agents was made independently in the Lauterbur and Leigh laboratories (Mendoca Dias and Lauterbur, 1986; Renshaw et al., 1986). Renshaw working with Leigh demonstrated that antibodies could be coupled to iron oxide particles for targeted MRI contrast and Lauterbur was interested in the ability to track cells in liver, spleen, and brain with iron oxide particles (Hawrylak et al., 1993; Mendoca Dias and Lauterbur, 1986). Cell tracking using iron oxide contrast has grown into a robust area of investigation in MRI (Bulte, 2009; Ho and Hitchens, 2004). Magnetic resonance was also beginning to be integrated with emerging transgenic mouse technology to study problems in physiology (Koretsky, 1995). Early papers demonstrated that expression of creatine kinase could be used as a reporter strategy by detecting the metabolic product phosphocreatine with magnetic resonance in *E. coli* (Koretsky and Traxler, 1989), yeast (Brindle et al., 1990) and liver in transgenic mice (Koretsky et al., 1990). Developing reporter protein strategies that enable detection of gene expression with MRI is now an active research area (Gilad et al., 2007a, 2007b). Measurement of gene expression of immediate early genes such as *cfos* has a long history of use for studying brain function (Barth, 2007), and MRI has the potential to do this non-invasively. Thus, prior to the first human BOLD fMRI results there was a body of work aimed at using MRI to monitor brain function including metabolic correlates of brain function, oxygenation, blood flow, calcium concentrations, cell tracking and gene expression. Indeed, these studies helped lay the foundation for the area we now call molecular imaging with MRI. Despite the success of BOLD fMRI there continues to be work attempting to find MRI techniques to measure brain function more directly than relying on vascular responses to infer neural activity. The goal of this contribution is to highlight some of the attempts to get closer to neurons using MRI with emphasis on those approaches that are now part of the toolbox of molecular imaging.

Direct measurement of membrane potential: the holy grail

Neurons require propagation of action potentials to transmit information. Direct measurement of membrane potential changes can be accomplished with patch clamp techniques (Nauen, 2011) and fast responding, membrane potential sensing optical dyes (Chavane et al., 2011; Peterka et al., 2011; Waggoner, 1979). Patch clamp techniques are limited to a small number of sites and, although widely used in brain slice work, remain a challenge to perform in vivo (Petersen, 2009). Optical membrane potential dyes have a small dynamic range and can only be imaged from the surface of the brain. Exciting new developments in fluorescent proteins that sense membrane potential look promising to enable two-photon fluorescence techniques to make multi-site measurement of membrane potential from deeper tissue and allow targeting to specific cell types (Kralj et al., 2011). Other electrophysiological tools such as MEG, EEG and extracellular electrodes measure extracellular currents or potentials generated by action potentials and are thus less direct than monitoring membrane potential changes. A MRI technique that can either directly measure membrane potentials or the effects of action potential propagation would enable more direct monitoring of neural function than is possible with BOLD.

There is a literature that attempts to measure the effects of fluctuating magnetic fields generated from action potential propagation without the need to resort to any contrast agents. There have been attempts to measure the effects of fluctuating magnetic fields associated with neural activity on the phase or apparent relaxation of MRI signals (Bandettini et al., 2005). No attempt will be made to summarize this field. It remains an active area of investigation but none of

the proposed measurements has yet proven to be widely adopted. There are also recent attempts to monitor the swelling of cells known for many years to occur during action potential propagation using diffusion based MRI techniques (Le Bihan, 2007). This also remains an active area of investigation but with no consensus yet on whether such approaches will be broadly applicable. The prediction that there is a significant Lorentz force on neurons due to propagating action potentials in a magnetic field causing small motion of neurons that may be detected by MRI has also been considered with contradictory conclusions at this stage (Song and Takahashi, 2001; Roth and Basser, 2009).

A number of approaches to making MRI agents directly sensitive to membrane potential can be envisioned. MRI contrast agents with properties similar to the optical membrane potential sensitive dyes can be imagined. Indeed, lipophilic cations have long been used to measure slow membrane potential changes using fluorescence with particular emphasis on mitochondrial membrane potential (Waggoner, 1979). There is a large literature in nuclear medicine using radiolabeled derivatives of lipophilic cations to measure delivery of agent but the fact that they distribute intracellularly based on membrane potential is not exploited (Liu, 2007; Madar et al., 2006). The membrane permeant triphenyl phosphonium ion has been used to measure slow potential changes and can be directly detected with ^{31}P NMR, although this has not been used in MRI. A complex of a DO3A-conjugated triphenylphosphonium cation with a paramagnetic metal chelated has been reported (Yang et al., 2007). However, to the best of my knowledge no work using MRI has yet been done in this direction to use these agents to attempt to measure neural potential changes.

More attractive for studying neural activity is the fast responding class of membrane potential optical dyes (Chavane et al., 2011; Peterka et al., 2011; Waggoner, 1979). A new class of fluorescent based protein indicators of fast membrane potential is being developed that show much promise (Kralj et al., 2011). The ideas used may be amenable to adaptation to an MRI based sensor. However, little work has been done to develop an MRI agent that can monitor fast membrane potentials. Paul Lauterbur's group did propose an idea aimed at measuring membrane potentials (Frank and Lauterbur, 1993). They embedded iron oxide particles in a gel matrix that would be expected to shrink and swell due to changes in electric field. Altering the distribution of particles caused changes in T_2 enabling a probe of membrane potential. This class of agent is based on T_2 changes allowing potential changes to be measured on fast time scales. Challenges of this approach are to find ways to embed the particles in the portion of the membrane where potentials are actually changing, as well as to make gels that are specific for membrane potential changes and that are not affected by other changes such as pH or Ca^{2+} . The idea to control the aggregation of iron oxide particles to make sensors has grown (Atanasijevic et al., 2006; Colomb et al., 2011; Haun et al., 2011; Perez et al., 2002), however, no further work on sensing membrane potential has been performed. While there would be many uses of MRI based ways to monitor membrane potential changes directly, it remains a holy grail in need of new maps.

Calcium dynamics using MRI

Measurement of intracellular calcium

When a neurotransmitter binds to its receptor on the postsynaptic neuron and triggers an action potential, Ca^{2+} enters the neuron eventually leading to release of neurotransmitter at the neuron's pre-synaptic terminal. Thus, measurement of Ca^{2+} influx or changes in intracellular Ca^{2+} concentrations is an excellent surrogate for membrane potential changes. As mentioned in the introduction, the development of robust fluorescent indicators of Ca^{2+} (Tsien, 1980)

and the recent generation of fluorescent proteins sensitive to Ca^{2+} (Looger and Griesbeck, 2012; Miyawaki et al., 1997) have been very important for studies of neural function in brain slices and in animal brains.

There have been a number of MRI based agents to measure Ca^{2+} . Based on the first generation of fluorescent Ca^{2+} indicators, ^{19}F based Ca^{2+} indicators were developed and applied to brain slice work (Bador-Goffer et al., 1990; Smith et al., 1983). It was demonstrated that a chelator with a well positioned ^{19}F could have a large chemical shift upon binding Ca^{2+} . The large concentrations needed compared to the free Ca^{2+} concentration raised concerns about buffering and techniques to deliver the agent efficiently to the intact brain have hampered work with these agents. There is now a growing literature developing Gadolinium (Gd) chelates as Ca^{2+} sensors (Angelovski et al., 2008; Li et al., 1999; Li et al., 2002; Mamedov et al., 2011; Hennig et al., 2011). The basic idea is to use a dual chelator one to hold the Gd and another to bind Ca^{2+} . The trick is to engineer the molecule so that water exposure to Gd in the chelate changes when Ca^{2+} binds. This enables changes in T_1 relaxivity due to changing Ca^{2+} concentrations by this class of agent. Ca^{2+} binding can vary accessibility to the first or second coordination sphere of water or change the exchange rate in order to alter T_1 relaxivity. Proof of principle, a growing list of possible agents, details of the mechanism for relaxivity changes, and potential problems such as aggregation have been studied (Li et al. 2002; Hennig et al., 2011; Mamedov et al., 2011; Mishra et al., 2011). There has also been the suggestion to use Ca^{2+} induced oligomerization of Gd dota-bisphosphonate conjugates as an indicator (Kubicek et al., 2010). Extremely large changes in relaxivity (200–500%) were reported on binding Ca^{2+} . Unfortunately this agent was not specific for Ca^{2+} over other divalent metal ions such as Mg^{2+} and Zn^{2+} . A problem with agents that change T_1 relaxivity is that typically calcium transients are very rapid (~10 ms) inside cells to keep up with spiking neurons. It is not clear how agents that change relaxivity on the T_1 time scale (~1 s) can keep up. However, as was the case with many of the early generation fluorescent indicators, the slow responding MRI agents may be useful for monitoring average calcium changes from a series of many action potentials. The trick will be to engineer the affinity and the off rates to enable this type of integration of calcium transients.

Rather than use changes in T_1 relaxivity, there has been some initial work using chemical exchange dependent saturation transfer (CEST agents) to monitor Ca^{2+} (Angelovski et al., 2011). CEST agents rely on measuring chemical exchange between a chemically shifted resonance on the agent and water (Ward et al., 2000). Rather than the fast exchange that Gd based agents rely on, these agents rely on slow chemical exchange between a proton on the CEST agent and water protons. Development of CEST agents made a major step forward when it was demonstrated that binding of water to a paramagnetic ion on a chelate could be used to induce a large chemical shift and have exchange rates useful for detection via CEST strategies. This class of agents is known as paraCEST agents (Zhang et al., 2001). There are now a growing number of CEST agents that measure enzymes, metabolites, zinc, pH and temperature (Viswanathan et al., 2010). A major advantage of CEST agents is that the exchange effect can be readily turned on and off with appropriate MRI sequences. The Ca^{2+} paraCEST agent relied on synthesizing calcium chelators to the paraCEST agents and optimizing for changes induced by changing Ca^{2+} (Angelovski et al., 2011). The chemical exchange effect of CEST agents builds up on a T_1 time scale and so these agents will be limited to following average Ca^{2+} levels.

There has not been any work in neural preparations using either the T_1 relaxivity based agents or the paraCEST based agents. An interesting suggestion is to use these paramagnetic based agents to monitor changes in extracellular calcium with neural activity (Angelovski et al., 2008; Mishra et al., 2011). Decreases in extracellular Ca^{2+} as large as 0.2 mM have been reported in brain slice

work due to influx of Ca^{2+} into cells during neural activity (Fedirko et al., 2007). An extracellular MRI agent would need to have low affinity for Ca^{2+} to respond to the changes in the mM levels of extracellular Ca^{2+} , making it difficult to engineer specificity for Ca^{2+} over other ions. However, if successful this would bypass problems with loading the agents to the intracellular space and problems associated with the relatively high concentrations needed for MRI causing buffering of intracellular Ca^{2+} .

Building on the idea to use density or aggregation state of iron oxide particles as a sensor, the Jasanoff lab has engineered iron oxide particles to enable Ca^{2+} to control their aggregation state (Atanasijevic et al., 2006). This builds on similar ideas from Josephson and Weissleder who have developed a number of sensors by controlling the aggregation of iron oxide particles (Perez et al., 2002; Haun et al., 2011). The Ca^{2+} indicator used the protein calmodulin as the calcium sensor on one set of particles and a peptide known to bind only to Ca^{2+} bound calmodulin on the other set of particles. An increase in calcium causes the particles to aggregate and altered the apparent T_2 of water (Atanasijevic et al., 2006). It is possible to use site directed mutagenesis to optimize the properties of this indicator (Green et al., 2006). Problems with this approach include delivering agents to the cytoplasm and the need to set the concentration of agent just right to get desired T_2 changes. Assembly to the aggregated state can be quite slow even in vitro and it is not clear how much the cytoskeleton may further slow the process. There has been recent work modeling the processes that control MRI relaxivity and the aggregation of iron oxide nanoparticles including ferritin (Bennett et al., 2008; Matsumoto and Jasanoff, 2008; Shapiro et al., 2006). Shapiro et al. also discuss time resolution limitations associated with the entire range of MRI based Ca^{2+} agents (Shapiro et al., 2006).

There has been recent interest in altering the expression of ferritin as a gene expression reporter strategy (see discussion below; Cohen et al., 2005; Genove et al., 2005). It may be possible to engineer chimeras of ferritin with Ca^{2+} binding proteins to enable neurons to express their own calcium indicators as is done with Ca^{2+} sensitive fluorescent proteins (Looger and Griesbeck, 2012; Miyawaki et al., 1997). This strategy would be a great step forward for animal studies.

Manganese Enhanced MRI (MEMRI) as a surrogate for calcium influx

Mn^{2+} was the first paramagnetic ion used to alter contrast in MRI (Lauterbur, 1973). Early work demonstrated that it was transported intracellularly, however, toxicity concerns hindered its development. With the advent of more sensitive MRI scanners the use of Mn^{2+} as a useful contrast agent has had a renaissance especially for brain imaging in animal models. Mn^{2+} is an essential heavy metal for life and as such there are many interesting biological processes to transport and sequester Mn^{2+} . One such mechanism is that Mn^{2+} can enter cells on voltage gated calcium channels. Therefore, the influx of Mn^{2+} into excitable tissue is a surrogate for Ca^{2+} influx. This idea was used to make functional maps of somatosensory areas in brain by Lin et al. in the rat brain (Lin and Koretsky, 1997). There is now a significant body of work that has used Mn^{2+} as a versatile contrast agent to study the brain (Boretius and Frahm, 2011; Inoue et al., 2011). There have been a few studies that have demonstrated excellent agreement between MEMRI maps and fMRI maps of neural representations (see Silva in this issue of NeuroImage). Recent work in the olfactory bulb (Chuang et al., 2009a), hypothalamus (Kuo et al., 2006; Parkinson et al., 2009), retina (Berkowitz et al., 2006; Berkowitz et al., 2009), motor cortex (Eschenko et al., 2010a, 2010b), nociceptive pathways (Yang et al., 2011), and the auditory midbrain in mice (Yu et al. 2005; Yu et al. 2007) demonstrates that MEMRI has specificity to image small neural representations as well as laminar activity and even an individual olfactory glomerulus. Many of the areas mapped would be too small for BOLD based fMRI

studies. For most areas of the brain, Mn^{2+} entry is rate limiting and so requires breaking the blood brain barrier (BBB). Novel approaches using focused ultrasound and BBB breaking antibodies have been used with MEMRI (Howles et al., 2010; Lu et al., 2010).

Mn^{2+} has been used to monitor activity in a few brain areas without breaking the BBB in areas that transport Mn^{2+} relatively quickly from the blood, such as the hypothalamus (Kuo et al., 2006) and the retina (Berkowitz et al., 2006). Another approach to avoid breaking the BBB has been to use very long periods of stimulation in the presence of Mn^{2+} (Eschenko et al., 2010a, 2010b; Yu et al., 2005). There are a number of limitations to using MEMRI. Mn^{2+} leaves the brain on the time scale of weeks (Chuang et al., 2009b) limiting its applicability. It is not yet clear what are the minimum number of action potentials required to enable enough accumulation of Mn^{2+} to enable detection but it is likely that many are required. Nor is it clear what the distribution of Mn^{2+} is between neurons and glial, since Mn^{2+} can accumulate in both cell types in an active brain area. The slow eflux enables MEMRI to make very high resolution maps and allows brain areas to be stimulated outside the magnet of MRI systems and imaged (Eschenko et al., 2010a, 2010b; Lin and Koretsky, 1997; Yu et al., 2005). The idea to use Mn^{2+} accumulation to map active brain areas has not yet been attempted in the human brain. Access through the BBB and concerns about toxicity are the main barriers (Eschenko et al., 2010a, 2010b; Jackson et al., 2011). However, work demonstrates that the FDA approved Mn^{2+} based contrast, MnDPDP, can deliver Mn^{2+} to the pituitary and choroid plexus (Wang et al., 1997), offering the intriguing possibility of using MEMRI to study human brain function.

Two other unique properties of Mn^{2+} are finding widespread use in animal models. Mn^{2+} is transported in brain in an anterograde enabling direct monitoring of brain connectivity with laminar specificity (Pautler et al., 1998; Tucciarone et al. 2009; Inoue et al., 2011; Zhang et al., 2010). The tracing occurs on a much slower time scale than uptake used to map neural activity. The second useful property of Mn^{2+} is that once accumulated in the brain, Mn^{2+} gives contrast that enables cytoarchitectural information to be obtained from MRI (Aoki et al., 2004; Bock et al., 2009; Silva et al., 2008; Watanabe et al., 2001). Indeed structures as small as individual olfactory glomeruli have been resolved using the different types of MEMRI experiments (Chuang et al., 2009a; Chuang et al., 2010).

MRI agents to measure neurotransmitter release

Another area that would enable MRI to get closer to neurons is monitoring neurotransmitter release. Spectroscopic techniques have enabled average glutamate recycling and GABA recycling rates to be measured (Novotny et al., 2003). While very useful for studying pathophysiology of the brain, the time scale of the experiments and the resolution do not lend themselves readily to monitoring functional brain states. Spectroscopic detection of GABA levels has been associated with neural activity and may represent changes in the “inhibitory” state of a brain area (Chen et al., 2005; Muthukumaraswamy et al., 2009).

Recently there has been exciting work using advanced laboratory protein evolution techniques to make a MRI sensor of dopamine (Shapiro et al., 2010). The idea was to use a heme based protein whose ligand binding alters water access to the metal in the heme. In this case bacterial cytochrome P450-BM3, which has a paramagnetic iron in its heme site and is accessible to water in its unliganded state was used. Addition of the natural ligand arachidonic acid to BM3 led to a decrease in T_1 relaxivity. Directed evolution was used and mutants with an increased affinity to dopamine were selected for further rounds of mutagenesis. After a number of rounds of mutagenesis, a protein that had high affinity for dopamine rather than arachidonic acid was produced creating an MRI sensor of dopamine. Preliminary results indicated that this agent will be useful for monitoring

extracellular dopamine in the rat brain (Shapiro et al., 2010). Optimization of the metal binding has enabled increasing the dynamic range (Lelyveld et al. 2011). Of course, one can imagine evolving protein sensors for other neurotransmitters and the resulting protein can be expressed and secreted by the neurons themselves, bypassing delivery issues. It will be exciting to see if these uses of advanced protein evolution approaches can be widely adapted for making agents suitable for studies in animals. Application to the human brain will have to await toxicity studies and efficient ways to deliver these proenzyme constructs through the BBB. An advantage is that these agents detect extracellular levels of neurotransmitter so there is no need to deliver the agents to the intracellular space.

Rather than monitor neurotransmitters directly it might be useful to monitor other constituents of neurotransmitter containing vesicles. In particular, many neurotransmitter vesicles are enriched in zinc. There have been a number of MRI agents designed to measure zinc. These are primarily based on approaches similar to those used for calcium sensitive agents. T_1 paramagnetic chelates or paraCEST chelates have been modified to contain zinc chelating groups that modulate the relaxivity or exchange upon binding zinc (Lee et al., 2010; Major et al., 2007; Mishra et al., 2011; Trowski et al., 2005; Zhang et al., 2007). As with the calcium indicators these have been primarily proof of principle studies. Recently, a MRI based zinc indicator has been used to measure extracellular zinc release from the pancreas in rodent in response to elevated glucose (Lubag et al., 2011). Insulin vesicles are high in zinc and the ability to have a surrogate marker of insulin release would be very important for following the progression of diabetes. In this case, a Gd based chelator (Gd DOTA-diBPEN) binds zinc which in turn imparts a high affinity of the chelate for albumin. The binding of the low molecular weight Gd chelate to albumin increases the T_1 relaxivity due to a change in correlation time (Esqueda et al., 2009). It would be very exciting if this approach proved to be sensitive enough to extend to monitoring release of zinc from central nervous system synapses!

MRI reporters of gene expression

A number of genes increase expression in response to neural activity. Monitoring the expression of these genes has been a surrogate for monitoring neural activity. Most important has been *cfos* and other early neural response genes such as *Arc* (Barth, 2007). The advent of fluorescent proteins has enabled rodent models that enable direct monitoring of increased gene expression due to brain activity (Barth, 2007; Wang et al., 2006). A similar strategy could be accomplished with MRI. There has been a long hunt for a robust MRI reporter strategy for monitoring gene expression. All gene reporter strategies rely on detecting the expression of a detectable protein or RNA. The MRI approaches all rely on expressing a protein that can alter magnetic resonance properties. Starting with work using creatine kinase (and subsequently the related arginine kinase) to alter metabolites that could be detected by magnetic resonance (Auricchio et al., 2001; Brindle et al., 1990; Koretsky and Traxler, 1989; Koretsky et al., 1990; Landis et al., 2006; Li et al., 2005; Walter et al. 2000), there has now been development of a large number of strategies (Gilad et al., 2007a, 2007b).

The enzyme beta-galactosidase has been used since the birth of molecular biology as a reporter protein and so it has been a target for development of a number of MRI agents that can detect this enzyme. A Gd chelate where the Gd is protected from water until the enzyme beta-galactosidase is expressed to cleave the chelate and cause an increase in T_1 relaxivity has been demonstrated (Louie et al., 2000). A number of similar Gd chelate strategies have now been reported (Chang et al., 2007; Hanaoka et al., 2008; Keliris et al., 2011). Shortcomings have been the relatively small change in relaxivity associated with enzymatic activation of these agents as well as the problem of distinguishing the amount of agent from the

amount of activation. A recent study reports on a Gd chelate that was designed to polymerize in the presence of beta-galactosidase and melanin (Arená et al., 2011). Fluorinated substrates have been designed so that the fluorine chemical shift changes when the substrate is cleaved by beta-galactosidase enabling ^{19}F NMR to detect gene expression (Cui et al., 2004; Liu et al., 2007). These agents are limited by the need for very high concentrations. Colorimetric agents have been used for functional assays of beta-galactosidase for a long time in isolated cells which rely on enzymatic activity to form a precipitate which is high in iron. Recently it has been demonstrated that these agents can be used in vivo for MRI (Bengtsson et al., 2010; Cui et al., 2010). In order to apply any of the gene expression reporter protein strategies that require exogenous contrast agent to the brain will require them to be permeable through the BBB.

Approaches to increase stored iron in a cell have the potential to be the most sensitive by altering T_2 relaxivity. There have been a number of proteins whose expression was controlled in an attempt to increase cellular iron, such as expressing the transferrin receptor that increases iron transport into cells (Koretsky et al., 1996). Increasing tyrosinase to produce melanin which binds iron has also been proposed as a potential reporter protein strategy to increase cellular iron (Weissleder et al., 1997). There has been growing interest in expressing ferritin, which is an iron storage protein. Changes in T_2 have been reported in mouse models that overexpress ferritin (Cohen et al., 2005; Cohen et al., 2007; Choi et al., 2012; Genove et al., 2005; Ziv et al., 2010). Ferritin has low MRI relaxivity and so large levels of expression are required. Dual expression of both transferrin receptor and ferritin has been reported, but cells needed to be pre-incubated with high iron to get significant changes in relaxivity compared to controls (Deans et al., 2006). Recent work has reported only small changes in MRI signals due to ferritin overexpression (Aung et al., 2009; Kim et al., 2010; Ono et al., 2009; Van de Velde et al., 2011). A transgenic mouse expressing human ferritin heavy chain in brain did not accumulate iron indicating that very high levels of expression may be required (Hasegawa et al., 2011). The high levels of expression and small signal changes detected with expression of the transferrin receptor or ferritin indicate that the idea to cause iron accumulation in cells via gene expression needs to be optimized. Recently, a more optimal ferritin with higher iron loading and a higher T_2 relaxivity has been reported and used to track neural precursors in the rodent brain (Iordanova et al., 2010; Iordanova and Ehrens, 2012). There is a large potential for optimizing ferritin since similar sized iron oxide particles have about 100 times higher relaxivity. Maybe the advanced protein evolution techniques used to make the MRI dopamine sensor can be applied successfully to improve ferritin. In addition to using the normal mammalian mechanisms to increase cell iron, there is some evidence that expression of the bacterial iron transporter MagA may be sufficient for producing magnetic nanoparticles in mammalian cells making it a potential reporter protein (Zurkiya et al., 2008; Goldhawk et al., 2009).

Other approaches to altering MRI contrast via expression of a gene have involved expressing peptides that enable efficient exchange of magnetization from lysine or arginine protons to water protons and applying CEST techniques (Gilad et al., 2007a, 2007b, Liu et al., 2011, 2012). Expressing cell surface proteins such as the transferrin receptor or the major histocompatibility class 1 antigen that can be targeted with antibodies carrying exogenous MRI contrast is yet another strategy that has been demonstrated (Moore et al., 1998; So et al., 2005). Expressing enzymes that can degrade the coating on iron oxide particles taken up in the endosomal pathway and leading to loss of MRI contrast has been suggested to detect gene expression with MRI (Granot and Shapiro, 2011). It should be possible to make manganese based agents that will increase contrast when degraded by specific enzymes (Shapiro and Koretsky, 2008). Finally, there has been work that targets MRI

contrast directly to expressed RNA (Liu et al., 2009; Liu et al., 2011, 2012). Oligonucleotides specific for transcribed RNAs from a number of interesting brain genes were complexed to iron oxide particles which are retained longer than control iron oxide particles. Since iron oxide particles usually are taken up into lysosomes, the added oligonucleotide probes must help the particle enter the cytoplasm in order to be able to hybridize to the target RNA. If similar results can be achieved by other groups this would represent an exciting approach to monitor gene expression.

A large variety of novel approaches are being developed to try and find a sensitive and specific approach that can be used as a gene expression reporter for MRI. In most of the cases demonstrated to date very strong gene expression promoters have been used to get enough protein to lead to detectable MRI signal or cells were first transfected to produce high levels of the reporter protein in vitro prior to transplanting into an animal. Even under these optimal circumstances the contrast changes reported have been small. The majority of the studies have been proof of principle with few applications yet. Thus, it is not clear that any of these strategies can be robustly used to monitor changes in normal cellular promoters such as found on the genes that respond to neural activity. It is also difficult to envision how these techniques will translate to work on the human brain. However, when a widely applicable approach is developed it will be very useful for rodent and non-human primate studies. The important use of the multiple colors available from fluorescent proteins makes it likely that different MRI reporter strategies that make use of T_1 , T_2 and CEST contrast will be important.

Conclusion

In the early 1990s, I had the good fortune of having an excellent graduate student at Carnegie Mellon University, Yi-Jen Lin. Yi-Jen joined my lab and we were worried that the new MRI hemodynamic based strategies for mapping the brain might have limited utility. This led to a hunt for alternative strategies and the development of MEMRI (Lin and Koretsky, 1997) as well as the thought to make gene expression reporters that might eventually be used for mapping brain activity (Koretsky et al., 1996). On the 20th anniversary of BOLD fMRI in humans, it has become clear that there was no reason for Yi-Jen and I to be worried! BOLD has developed beyond anyone's wildest imagination to study all aspects of brain function, open up the field of resting state connectivity, and decode what is on our minds! This has raised the bar for any new MRI tools that hope to measure brain function.

Despite this tremendous success of BOLD fMRI, there remains a very active field of inquiry aiming to develop approaches to move MRI closer to directly measuring neural activity. The dependence of fMRI on vascular responses still lead to ambiguities related to localization of neural activity, timing of neural activity, defining the neurotransmitter systems responsible for neural activity, and defining the cellular source of the neural activity. The dream of molecular imaging techniques would be to add specificity to fMRI studies that would circumvent the problems associated with relying on vascular responses to detect neural responses indirectly. There has been some progress and a number of very interesting ways to control MRI contrast to sense a growing list of biological molecules and biological processes. The impact so far has been in defining new modes of MRI contrast and studying brain function in animal models. In this growing field of molecular imaging, animal studies are where the exciting MRI development originates, as was the case for BOLD fMRI in 1990 (Ogawa et al., 1990a, Ogawa and Lee, 1990, Ogawa et al., 1990b). Translation of these new molecular imaging strategies to the human brain seems a long way off, but our hope is that when we look back twenty years from now a range of MRI approaches will be available to get us closer to the neurons.

Acknowledgments

Alan Koretsky is supported by the intramural research program of the National Institute of Neurological Disorders and Stroke, NIH. The author acknowledges the kind invitation from Peter Bandettini to write this article and thanks Afonso Silva for encouragement and comments.

References

- Ackerman, J.J., Ewy, C.S., Kim, S.G., Shalwitz, R.A., 1987. Deuterium magnetic resonance in vivo: the measurement of blood flow and tissue perfusion. *Ann. NY. Acad. Sci.* 508, 89–98.
- Angelovski, G., Fuskova, P., Mamedov, I., Canals, S., Toth, E., Logothetis, N.K., 2008. Smart Magnetic Resonance Imaging Agents that sense extracellular calcium fluctuations. *ChemBioChem* 9, 1729–1734.
- Angelovski, G., Chauvin, T., Pohmann, R., Logothetis, N.K., Toth, E., 2011. Calcium responsive paramagnetic CEST agents. *Biorg. Med. Chem.* 19, 1097–1105.
- Aoki, I., Wu, Y.J., Silva, A.C., Lynch, R.M., Koretsky, A.P., 2004. In vivo detection of neuroarchitecture in the rodent brain using manganese enhanced MRI. *NeuroImage* 22, 1046–1059.
- Arai, T., Nakao, S., Mori, K., Ishimori, K., Morishima, I., Miyazawa, T., Fritz-Zieroth, B., 1990. Cerebral oxygen utilization analyzed by the use of oxygen-17 and its nuclear magnetic resonance. *Biochem. Biophys. Res. Commun.* 169, 153–158.
- Arena, F., Singh, J.B., Gianolio, E., Stefania, R., Aime, S., 2011. b-Gal gene expression MRI reporter in melanoma tumor cells. Design, synthesis, and in vitro and in vivo testing of a Gd(III) containing probe forming a high relaxivity, melanin-like structure upon b-Gal enzymatic activation. *Bioconjug. Chem.* 22, 2625–2635.
- Atanasijevic, T., Shusteff, M., Fam, P., Jasanoff, A., 2006. Calcium sensitive MRI contrast agents based on superparamagnetic iron oxide nanoparticles and calmodulin. *Proc. Natl. Acad. Sci.* 103, 14707–14712.
- Aung, W., Hasegawa, S., Koshikawa-Yana, M., Obata, T., Ikehira, H., Furukawa, T., Aoki, I., Saga, T., 2009. Visualization of in vivo electroporation-mediated transgene expression in experimental tumors by optical and magnetic resonance imaging. *Gene Ther.* 16, 830–839.
- Auricchio, A., Zhou, R., Wilson, J.M., Glickson, J.D., 2001. In vivo detection of gene expression in liver by ³¹P nuclear magnetic resonance spectroscopy employing creatine kinase as a marker gene. *Proc. Natl. Acad. Sci. USA* 98, 5205–5210.
- Bador-Goffer, R.S., Ben-Yoseph, O., Dolin, S.J., Morris, P.G., Smith, G.A., Bachelard, H.S., 1990. Use of 1,2-bis(2-amino-5-fluorophenoxy)ethane-N, N, N', N'-tetraacetic acid (5FBAPTA) in the measurement of free intracellular calcium in the brain by ¹⁹F-nuclear magnetic resonance spectroscopy. *J. Neurochem.* 55, 878–884.
- Bandettini, P.A., Petridou, N., Bodurka, J., 2005. Direct detection of neuronal activity with MRI: fantasy, possibility, or reality? *Appl. Magn. Reson.* 29, 65–75.
- Barth, A., 2007. Visualizing circuits and systems using transgenic reporters of neural activity. *Curr. Opin. Neurobiol.* 17 (567), 571.
- Bengtsson, N.E., Brown, G., Scott, E.W., Walter, G.A., 2010. lacZ as a genetic reporter for real time MRI. *Magn. Reson. Med.* 63, 745–753.
- Bennett, K.M., Shapiro, E.M., Sotak, C.H., Koretsky, A.P., 2008. Controlled aggregation of ferritin to modulate MRI relaxivity. *Biophys. J.* 95, 342–351.
- Berkowitz, B.A., Roberts, R., Goebel, D.J., Luan, H., 2006. Noninvasive and simultaneous imaging of layer-specific retinal functional adaptation by manganese-enhanced MRI. *Invest. Ophthalmol. Vis. Sci.* 47, 2668–2674.
- Berkowitz, B.A., Roberts, R., Oleske, D.A., Chang, M., Schafer, S., Bissig, D., Gradianu, M., 2009. Quantitative mapping of ion channel regulation by visual cycle activity in rodent photoreceptors in vivo. *Invest. Ophthalmol. Vis. Sci.* 50, 1880–1885.
- Bock, N.A., Kocharyan, A., Silva, A.C., 2009. Manganese enhanced MRI visualizes V1 in the non-human primate visual cortex. *NMR Biomed.* 22, 730–736.
- Boretius, S., Frahm, J., 2011. Manganese-enhanced magnetic resonance imaging. *Methods Mol. Biol.* 771, 531–568.
- Brindle, K., Braddock, P., Fulton, S., 1990. ³¹P NMR measurements of the ADP concentration in yeast cells genetically modified to express creatine kinase. *Biochemistry* 29, 3295–3302.
- Bulte, J.W., 2009. In vivo MRI cell tracking: clinical studies. *AJR Am. J. Roentgenol.* 193, 314–325.
- Chang, Y.T., Cheng, C.M., Su, Y.Z., Lee, W.T., Hsu, J.S., Liu, G.C., Cheng, T.L., Wang, Y.M., 2007. Synthesis and characterization of a new bioactivated paramagnetic gadolinium(III) complex [Gd(DOTA-FPG)(H₂O)] for tracing gene expression. *Bioconjug Chem.* 18, 1716–1727.
- Chavane, F., Sharon, D., Jancke, D., Marre, O., Fregnac, Y., Grinvald, A., 2011. Lateral spread of orientation selectivity in V1 is controlled by intracortical cooperativity. *Front. Syst. Neurosci.* 5, 4 Epub ahead of print.
- Chen, Z.G., Silva, A.C., Yang, J.H., Shen, J., 2005. Elevated endogenous GABA levels correlates with decreased fMRI signals in the rat brain during acute inhibition of GABA transaminase. *J. Neurosci. Res.* 79, 383–391.
- Choi, S.H., Cho, H.R., Kim, H.S., Kim, Y.H., Kang, K.W., Kim, H., Moon, W.K., 2012. Imaging and quantification of metastatic melanoma cells in lymph nodes with a ferritin MR reporter in living mice. *NMR Biomed.* 25, 737–745.
- Chuang, K.H., Lee, J.H., Silva, A.C., Belluscio, L., Koretsky, A.P., 2009a. Manganese enhanced MRI reveals functional circuitry in response to odorant stimuli. *NeuroImage* 44, 363–372.
- Chuang, K.H., Koretsky, A.P., Sotak, C.H., 2009b. Temporal changes in the T1 and T2 relaxation rates (Delta R1 and Delta R2) in the rat brain are consistent with the tissue-clearance rates of elemental manganese. *Magn. Reson. Med.* 61, 1528–1532.
- Chuang, K.H., Belluscio, L., Koretsky, A.P., 2010. In vivo detection of individual glomeruli in the rodent olfactory bulb using manganese enhanced MRI. *NeuroImage* 49, 1350–1356.
- Clark Jr., L.C., Ackerman, J.L., Thomas, S.R., Millard, R.W., Hoffman, R.E., Pratt, R.G., Ragle-Cole, H., Kinsey, R.A., Janakiraman, R., 1984. Perfluorinated organic liquids and emulsions as biocompatible NMR imaging agents for ¹⁹F and dissolved oxygen. *Adv. Exp. Med. Biol.* 180, 835–845.
- Cohen, B., Daffni, H., Meir, G., Harmelin, A., Neeman, M., 2005. Ferritin as an Endogenous MRI Reporter for Noninvasive Imaging of Gene Expression in C6 Glioma Tumors. *J. Neurosci.* 25, 109–117.
- Cohen, B., Ziv, K., Plaks, V., Israely, T., Kalchenko, V., Harmelin, A., Benjamin, L.E., Neeman, M., 2007. MRI detection of transcriptional regulation of gene expression in transgenic mice. *Nat. Med.* 13, 498–503.
- Colomb, J., Louie, K., Massia, S.P., Bennett, K.M., 2011. Self-degrading, MRI detectable hydrogel sensors with picomolar target sensitivity. *Magn. Reson. Med.* 64, 1792–1799.
- Cui, W., Otten, P., Li, Y., Koeneman, K.S., Yu, J., Mason, R.P., 2004. Novel NMR approach to assessing gene transfection: 4-fluoro-2-nitrophenyl-beta-D-galactopyranoside as a prototype reporter molecule for beta-galactosidase. *Magn. Reson. Med.* 51, 616–620.
- Cui, W., Liu, L., Kodibagker, V.D., Mason, R.P., 2010. S-Gal, a novel ¹H MRI reporter for beta-galactosidase. *Magn. Reson. Med.* 64, 65–71.
- Deans, A.E., Wadghiri, Y.Z., Bernas, L.M., Yu, X., Rutt, B.K., Turnball, D.H., 2006. Cellular MRI contrast via co-expression of transferrin receptor and ferritin. *Magn. Reson. Med.* 56, 51–59.
- Deuel, R.K., Yue, G.M., Sherman, W.R., Schickner, D.J., Ackerman, J.J., 1985. *Science* 228, 1329–1330.
- Eschenko, O., Canals, S., Simanova, I., Logothetis, N., 2010a. Behavioral, electrophysiological and histopathological consequences of systemic manganese administration in MEMRI. *Magn. Reson. Imaging* 28, 1165–1174.
- Eschenko, O., Canals, S., Simanova, I., Beyerlein, M., Murayama, Y., Logothetis, N.K., 2010b. Mapping of functional brain activity in freely behaving rats during voluntary running using manganese-enhanced MRI: implications for longitudinal studies. *NeuroImage* 49, 2544–2555.
- Esqueda, A.C., Lopez, J.A., Andreu-de-Riquer, G., Alvarado-Monzon, J.C., Ratnakar, J., Lubag, A.J., Sherry, A.D., De Leon-Rodriguez, L.M., 2009. A new gadolinium-based zinc sensor. *J. Am. Chem. Soc.* 131, 11387–11391.
- Fedirko, N., Avshalumov, M., Rice, M.E., Chesler, M., 2007. Regulation of postsynaptic Ca²⁺ influx in hippocampal CA1 pyramidal neurons via extracellular carbonic anhydrase. *J. Neurosci.* 27, 1167–1175.
- Frank, S., Lauterbur, P.C., 1993. Voltage sensitive magnetic gels as magnetic resonance monitoring agents. *Nature* 363, 334–336.
- Genove, G., DeMarco, U., Xu, H., Goins, W.F., Ahrens, E.T., 2005. A new transgene reporter for in vivo magnetic resonance imaging. *Nat. Med.* 11, 450–454.
- Gilad, A.A., McMahon, M.T., Walczak, P., Winnard Jr., P.T., Raman, V., van Laarhoven, H.W., Skoglund, C.N., Bulte, J.W., van Zijl, P.C., 2007a. Artificial reporter gene providing MRI contrast based on proton exchange. *Nat. Biotechnol.* 25, 217–219.
- Gilad, A.A., Winnard Jr., P.T., van Zijl, P.C., Bulte, J.W., 2007b. Developing MR reporter genes: promised and pitfalls. *NMR Biomed.* 20, 275–290.
- Goldhawk, D.E., Lemaire, C., McCreary, C.R., McGirr, R., Dhanvantari, S., Thompson, R.T., Figueredo, R., Koropatnick, J., Foster, P., Prato, F.S., 2009. Magnetic resonance imaging of cells overexpressing MagA, an endogenous contrast agent for live cell imaging. *Mol. Imaging* 8, 129–139.
- Granot, D., Shapiro, E.M., 2011. Release activation of iron oxide nanoparticles: (REACTION) a novel environmentally sensitive MRI paradigm. *Magn. Reson. Med.* 65, 1253–1259.
- Green, D.F., Dennis, A.T., Fam, P.S., Tidor, B., Jasanoff, A., 2006. Rational design of new binding specificity by simultaneous mutagenesis of calmodulin and a target peptide. *Biochemistry* 45, 12547–12559.
- Hanaoka, K., Kikuchi, K., Terai, T., Kmatsu, T., Nagano, T., 2008. A Gd³⁺-based magnetic resonance imaging contrast agent sensitive to beta-galactosidase activity utilizing a receptor-induced magnetization enhancement (RIME) phenomenon. *Chemistry* 14, 987–995.
- Hasegawa, S., Saito, S., Takanashi, J., Morokoshi, Y., Furukawa, T., Saga, T., Aoki, I., 2011. Evaluation of ferritin-overexpressing brain in newly developed transgenic mice. *Magn. Reson. Imaging* 29, 179–184.
- Haun, J.B., Yoon, T.J., Lee, H., Weissleder, R., 2011. Molecular detection of biomarkers and cells using magnetic nanoparticles and diagnostic magnetic resonance. *Methods Mol. Biol.* 726, 33–49.
- Hawrylak, N., Ghosh, P., Broadus, J., Schlueter, C., Greenough, W.T., Lauterbur, P.C., 1993. Nuclear magnetic resonance (NMR) imaging of iron oxide labeled neural transplants. *Exp. Neurol.* 121, 181–192.
- Hennig, J., Mamedov, I., Fouskova, P., Toth, E., Logothetis, N.K., Angelovski, G., Mayer, H.A., 2011. Influence of calcium induced aggregation on the sensitivity of aminobis(methylenephosphonate)-containing potential MRI contrast agents. *Inorg. Chem.* 50, 6472–6481.
- Ho, C., Hitchens, K., 2004. A non-invasive approach to detecting organ rejection by MRI: monitoring the accumulation of immune cells at the transplanted organ. *Curr. Pharm. Biotechnol.* 5, 551–566.
- Howles, G.P., Qi, Y., Johnson, G.A., 2010. Ultrasonic disruption of the blood brain barrier enables in vivo functional mapping of the mouse barrel field cortex with manganese-enhanced MRI. *NeuroImage* 50, 1464–1471.
- Inoue, T., Majid, T., Pautler, R.G., 2011. Manganese enhanced MRI (MEMRI): neurophysiological applications. *Rev. Neurosci.* Epub ahead of print.
- Iordanova, B., Ehrens, E.T., 2012. In vivo magnetic resonance imaging of ferritin-based reporter visualizes native neuroblast migration. *NeuroImage* 59, 1004–1012.

- Iordanova, B., Robison, C.S., Ahrens, E.T., 2010. Design and characterization of a chimeric ferritin with enhanced iron loading and transverse magnetic relaxation rate. *J. Biol. Inorg. Chem.* 15, 957–965.
- Jackson, S.J., Hussey, R., Jansen, M.A., Merrifield, G.D., Marshall, I., MacLulich, A., Yau, J.L., Bast, T., 2011. Manganese-enhanced magnetic resonance imaging (MEMRI) of rat brain after systemic administration of MnCl₂: hippocampal signal enhancement without disruption of hippocampus-dependent behavior. *Behav. Brain Res.* 216, 293–300.
- Keliris, A., Ziegler, T., Mishra, R., Pohmann, R., Sauer, M.G., Ugurbil, K., Engelmann, J., 2011. Synthesis and characterization of a cell-permeable bimodal contrast agent targeting β-galactosidase. *Bioorg. Med. Chem.* 19, 2529–2540.
- Kim, H.S., Cho, H.R., Choi, S.H., Woo, J.S., Moon, W.K., 2010. In vivo imaging of tumor transduced with bimodal lentiviral vector encoding human ferritin and green fluorescent protein on a 1.5T clinical magnetic resonance scanner. *Cancer Res.* 70, 7315–7324.
- Koretsky, A.P., 1995. Investigation of cell physiology in the animal using transgenic technology. *Am. J. Physiol.* 262, C261–C275.
- Koretsky, A.P., Traxler, B.A., 1989. The B isozyme of creatine kinase is active as a fusion protein in *Escherichia coli*: in vivo detection by ³¹P NMR. *FEBS Lett.* 243, 8–12.
- Koretsky, A.P., Brosnan, M.J., Chen, L.H., Chen, J.D., Van Dyke, T., 1990. NMR detection of the expression of creatine kinase in the liver of transgenic mice: determination of free ADP levels. *Proc. Natl. Acad. Sci. USA* 87, 3112–3116.
- Koretsky, A.P., Lin, Y.-J., Schorle, H., Jaenisch, 1996. Genetic control of MRI contrast by expression of the transferring receptor. *Proceedings of the International Society of Magn Reson Med.* 4, 69.
- Kralj, J.M., Hochbaum, D.R., Douglass, A.D., Cohen, A.E., 2011. Electrical spiking in *E. coli* probed with a fluorescent voltage-indicating protein. *Science* 333, 345–348.
- Kubicek, V., Vitha, T., Kotek, J., Hermann, P., Vander Elst, L., Muller, R.N., Lukes, I., Peters, J.A., 2010. Towards MRI contrast agents responsive to Ca(II) and Mg(II) ions: metal induced oligomerization of dota-bisphosphonate conjugates. *Contrast Media Mol. Imaging* 5, 294–296.
- Kuo, Y.T., Herlithy, A.H., So, P.W., Bell, J.D., 2006. Manganese-enhanced magnetic resonance imaging (MEMRI) without compromise of the blood–brain barrier detects hypothalamic neuronal activity in vivo. *NMR Biomed.* 19, 1028–1034.
- Landis, C.S., Yamanouchi, K., Zhou, H., Mohan, S., Roy-Chowdhury, N., Shafritz, D.A., Koretsky, A.P., Roy-Chowdhury, J., Hetherington, H.P., Guha, C., 2006. Noninvasive evaluation of liver repopulation by transplanted hepatocytes using ³¹P MRS imaging in mice. *Hepatology* 44, 1250–1258.
- Lauterbur, P., 1973. Image formation by induced local interactions: examples employing nuclear magnetic resonance. *Nature* 242, 190–191.
- Le Bihan, D., 2007. The “wet mind”: water and functional neuroimaging. *Phys. Med. Biol.* 52, R57–R90.
- Lee, T., Zhang, X.A., Dhar, S., Faas, H., Lippard, S.J., Jasanoff, A., 2010. In vivo imaging with a cell permeable porphyrin-based MRI contrast agent. *Chem. Biol.* 17, 665–673.
- Lelyveld, V.S., Brustad, E., Arnold, F.H., Jasanoff, A., 2011. Metal-substituted protein MRI contrast agents engineered for enhanced relaxivity and ligand sensitivity. *J. Am. Chem. Soc.* 133, 649–651.
- Li, W.H., Fraser, S.E., Meade, T.J., 1999. *J. Am. Chem. Soc.* 121, 1413–1414.
- Li, W.H., Parigi, G., Fragai, M., Luchinat, C., Meade, T.J., 2002. Mechanistic studies of a calcium dependent MRI contrast agent. *Inorg. Chem.* 41, 4018–4024.
- Li, Z., Qiao, H., Lebherz, C., Choi, S.R., Zhou, X., Gao, G., Kung, H.F., rader, D.J., Wilson, J.M., Glickson, J.D., Zhou, R., 2005. Creatine kinase a magnetic resonance-detectable marker gene for quantification of liver-directed gene transfer. *Hum Gene Ther.* 16, 1429–1438.
- Lin, Y.J., Koretsky, A.P., 1997. Manganese ion enhances T1-weighted MRI during brain activation: an approach to direct imaging of brain function. *Magn. Reson. Med.* 38, 378–388.
- Liu, S., 2007. Ether and crown ether-containing cationic ^{99m}Tc complexes useful as radiopharmaceuticals for heart imaging. *Dalton Trans.* 12, 1183–1193.
- Liu, L., Kodibagkar, V.D., Yu, J.X., Mason, R.P., 2007. 19F-NMR detection of lacZ gene expression via the enzymatic hydrolysis of 2-fluoro-4 nitrophenyl β-D-galactopyranoside in vivo in PC3 prostate tumor xenografts in the mouse. *FASEB J.* 21, 2014–2019.
- Liu, C.H., Ren, J.Q., Yang, J., Liu, C.M., Mandeville, J.B., Bhide, P.G., Yanagawa, Y., Liu, P.K., 2009. DNA-based MRI probes for specific detection of chronic exposure to amphetamine in living brains. *J. Neurosci.* 29, 10662–10670.
- Liu, G., Bulte, J.W., Gilad, A.A., 2011. CEST MRI reporter genes. *Methods Mol Biol.* 711, 271–280.
- Liu, C.H., Ren, J.Q., You, Z., Yang, J., Liu, C.M., Uppal, R., Liu, P.K., 2012. Noninvasive detection of neural progenitor cells in living brains by MRI. *FASEB J.* 26, 1652–1662.
- Looger, L.L., Griesbeck, O., 2012. Genetically encoded neural activity indicators. *Curr. Opin. Neurobiol.* 22, 18–23.
- Louie, A.Y., Huber, M.M., Ahrens, E.T., Rothbacher, U., Moats, R., Jacobs, R.E., Fraser, S.E., Meade, T.J., 2000. In vivo visualization of gene expression using magnetic resonance imaging. *Nat. Biotech.* 18, 321–325.
- Lu, H., Denny, S., Zuo, Y., Rea, W., Wang, L., Chefer, S.I., Vaupel, D.B., Yang, Y., Stein, E.A., 2010. Temporary disruption of the rat blood–brain barrier with a monoclonal antibody: a novel method for dynamic manganese-enhanced MRI. *NeuroImage* 50, 7–14.
- Lubag, A.J., De Leon-Rodriguez, L.M., Burgess, S.C., Sherry, A.D., 2011. Noninvasive MRI of b-cell function using a Zn²⁺-responsive agent. *Proc. Natl. Acad. Sci.* 108, 18400–18405.
- Madar, I., Ravert, H.T., Hilton, J., Volokh, L., Dannals, R.F., Frost, J.J., Hare, J.M., 2006. Characterization and uptake of the new PET imaging compound 18F-fluorobenzyl triphenyl phosphonium in dog myocardium. *J. Nucl. Med.* 47, 1359–1366.
- Major, J.L., Parigi, G., Luchinat, C., Meade, T.J., 2007. The synthesis and in vitro testing of a zinc-activated MRI contrast agent. *Proc. Natl. Acad. Sci.* 104, 13881–13886.
- Mamedov, I., Logothetis, N.K., Angelovski, G., 2011. Structure related variable responses of calcium sensitive MRI probes. *Org. Biomol. Chem.* 9, 5816–5824.
- Matsumoto, Y., Jasanoff, A., 2008. T2 relaxation induced by clusters of superparamagnetic nanoparticles: Monte Carlo simulations. *Magn Reson Imaging* 26, 994–998.
- Mendoca Dias, M.H., Lauterbur, P.C., 1986. Ferromagnetic particles as contrast agents for magnetic resonance imaging of liver and spleen. *Magn. Reson. Med.* 3, 328–330.
- Mishra, A., Logothetis, N.K., Parker, D., 2011. Critical in vitro evaluation of responsive MRI contrast agents for calcium and zinc. *Chemistry* 17, 1529–1537.
- Miyawaki, A., Llopis, J., Heim, R., McCaffery, J.M., Adams, J.A., Ikura, M., Tsien, R.Y., 1997. Fluorescent indicators for Ca²⁺ based on green fluorescent proteins and calmodulin. *Nature* 388, 882–887.
- Moore, A., Basilion, J.P., Chiocca, E.A., Weissleder, R., 1998. Measuring transferring receptor gene expression by NMR imaging. *Biochim. Biophys.* 1402, 239–249.
- Moseley, M.E., Cohen, Y., Mintorovich, J., Chileuitt, L., Shimizu, H., Kucharczyk, J., Wendland, M.F., Weinstein, P.R., 1990a. Early detection of regional ischemia in cats: comparison of diffusion- and T2-weighted MRI and spectroscopy. *Magn. Reson. Med.* 14, 330–346.
- Moseley, M.E., Cohen, Y., Kucharczyk, J., Mintorovitch, J., Asgari, H.S., Wendland, M.F., Tsuruda, J., Norman, D., 1990b. Diffusion-weighted MR imaging of anisotropic water diffusion in cat central nervous system. *Radiology* 176, 439–445.
- Muthukumaraswamy, S.D., Edden, R.A.E., Jones, D.K., Swettenham, J.B., Singh, K.D., 2009. Resting GABA concentration predicts peak gamma frequency and fMRI amplitude in response to visual stimulation in humans. *Proc. Natl. Acad. Sci. USA* 106, 8356–8361.
- Nauen, D.W., 2011. Methods of measuring activity at individual synapses: a review of techniques and the findings they have made possible. *J. Neurosci Methods* 194, 195–205.
- Novotny Jr., E.J., Fulbright, R.K., Pearl, P.L., Gibson, K.M., Rothman, D.L., 2003. Magnetic resonance spectroscopy of neurotransmitters in human brain. *Ann. Neurol.* 54 (Suppl. 6), S25–S31.
- Ogawa, S., Lee, T.M., 1990. Magnetic resonance imaging of blood vessels at high fields: in vivo and in vitro measurements and image simulation. *Magn. Reson. Med.* 16, 9–18.
- Ogawa, S., Lee, T.M., Kay, A.R., Tank, D.W., 1990a. Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proc. Natl. Acad. Sci. USA* 87, 9868–9872.
- Ogawa, S., Lee, T.M., Nayak, A., Glynn, P., 1990b. Oxygenation-sensitive contrast in magnetic resonance image of rodent brain at high magnetic fields. *Magn. Reson. Med.* 14, 68–78.
- Ono, K., Fuma, K., Tabata, K., Sawada, M., 2009. Ferritin reporter used for gene expression imaging by magnetic resonance. *Biochem. Biophys. Res. Comm.* 338, 589–594.
- Parkinson, J.R., Chaudhri, B.J., 2009. Imaging appetite-regulating pathways in the central nervous system using manganese-enhanced magnetic resonance imaging. *Neuroendocrinology* 89, 121–130.
- Pautler, R.G., Silva, A.C., Koretsky, A.P., 1998. In vivo neuronal tract tracing using manganese-enhanced magnetic resonance imaging. *Magn. Reson. Med.* 40, 740–748.
- Perez, J.M., O’Loughin, T., Simeone, F.J., Weissleder, R., Josephson, L., 2002. DNA-based magnetic nanoparticle assembly acts as a magnetic relaxation nanoswitch allowing screening of DNA-cleaving agents. *J. Am. Chem. Soc.* 124, 2856–2857.
- Peterka, D.S., Takahashi, H., Yuste, R., 2011. Imaging voltage in neurons. *Neuron* 69, 9–21.
- Petersen, C.C., 2009. Genetic manipulation, whole cell recordings and functional imaging of the sensorimotor cortex of behaving mice. *Acta Physiol.* 195, 91–99.
- Pritchard, J.W., Schulman, R.G., 1986. NMR Spectroscopy of brain metabolism in vivo. *Annu. Rev. Neurosci.* 9, 61–85.
- Radda, G.K., 1992. Control, bioenergetics, and adaptation in health and disease: noninvasive biochemistry from nuclear magnetic resonance. *FASEB J.* 6, 3032–3038.
- Renshaw, P.F., Owen, C.S., McLaughlin, A.C., Frey, T.G., Leigh Jr., J.S., 1986. Ferromagnetic contrast agents: a new approach. *Magn. Reson. Med.* 3, 217–225.
- Roth, B.J., Basser, P.J., 2009. Mechanical model of neural tissue displacement during Lorentz effect imaging. *Magn. Reson. Med.* 61, 59–64.
- Schnall, M.D., Bolinger, L., Renshaw, P.F., Haselgrove, J.C., Subramanian, V.H., Eleff, S.M., Barlow, C., Leigh Jr., J.S., Chance, B., 1987. Multinuclear MR imaging: a technique for combined anatomic and physiologic studies. *Radiology* 162, 863–866.
- Shapiro, E.M., Koretsky, A.P., 2008. Convertible manganese contrast for molecular and cellular MRI. *Magn. Reson. Med.* 60, 265–269.
- Shapiro, M.G., Atanasijevic, T., Faas, H., Westmeyer, G.G., Jasanoff, A., 2006. Dynamic imaging with MRI contrast agents: quantitative considerations. *Magn Reson. Imaging* 24, 449–462.
- Shapiro, M.G., Westmeyer, G.G., Romero, P.A., Szablowski, J.O., Kuster, B., Shah, A., Otey, C.R., Langer, R., Arnold, F.H., Jasanoff, A., 2010. Directed evolution of a magnetic resonance imaging contrast agent for noninvasive imaging of dopamine. *Nat. Biotech.* 28, 264–270.
- Silva, A.C., Lee, J.H., Wu, C.W., Tucciarone, J., Pelled, G., Aoki, I., Koretsky, A.P., 2008. Detection of cortical laminar architecture using manganese enhanced MRI. *J. Neurosci. Methods* 167, 246–257.
- Smith, G.A., Hesketh, R.T., Metcalfe, J.C., Feeney, J., Morris, P.G., 1983. Intracellular calcium measurements by 19F NMR of fluorine-labeled chelators. *Proc. Natl. Acad. Sci. U. S. A.* 80, 7178–7182.
- So, P.W., Hotee, S., Herlithy, A.H., Bell, J.D., 2005. Generic method for imaging transgene expression. *Magn. Reson. Med.* 54, 218–221.
- Song, A.W., Takahashi, A.M., 2001. Lorentz effect imaging. *Magn Reson Imaging* 19, 763–767.
- Trowski, R., Ren, J., Kalman, F.K., Sherry, A.D., 2005. Selective sensing of zinc ions with a paracetol contrast agent. *Angew. Chem. Int. Ed. Engl.* 44, 6920–6923.
- Tsien, R.Y., 1980. New calcium indicators and buffers with high selectivity against magnesium and protons: design, synthesis and properties of prototype structures. *Biochemistry* 19, 2396–2404.

- Tucciarone, J., Chuang, K.H., Dodd, S.J., Silva, A., Pelled, G., Koretsky, A.P., 2009. Layer specific tracing of corticocortical and thalamocortical connectivity in the rodent using manganese enhanced MRI. *NeuroImage* 44, 923–931.
- Van de Velde, G., Rangarajan, J.R., Toelen, J., Dresselaers, T., Ibrahim, A., Krylychkina, O., Vreys, F., Van der Linden, A., Maes, F., Debyser, Z., Himmelreich, U., Baekelandt, V., 2011. Evaluation of the specificity and sensitivity of ferritin as an MRI reporter gene in the mouse brain using lentiviral and adeno-associated viral vectors. *Gene Ther.* 18, 594–605.
- Viswanathan, S., Kovacs, Z., Green, K.N., Ratnakar, J., Sherry, A.D., 2010. Alternatives to Gadolinium-based metal chelates for magnetic resonance imaging. *Chem. Rev.* 110, 2960–3018.
- Waggoner, A.S., 1979. Dye indicators of membrane potential. *Annu. Rev. Biophys. Bioeng.* 8, 47–68.
- Walter, G., Barton, E.R., Sweeny, H.L., 2000. Noninvasive measurement of gene expression in skeletal muscle. *Proc. Natl. Acad. Sci. USA* 97, 5151–5155.
- Wang, C., Gordon, P.B., Hustvedt, S.O., Grant, D., Sterud, A.T., Martinsen, I., Ahlstrom, H., Hemmingsson, A., 1997. MR imaging properties and pharmacokinetics of MnDPDP in healthy volunteers. *Acta Radiol.* 38 (4 Pt 2), 665–676.
- Wang, K.H., Majewska, A., Schummers, J., Farley, B., Hu, C., Sur, M., Tonnegawa, S., 2006. In vivo two-photon imaging reveals a role of arc in enhancing orientation specificity in visual cortex. *Cell* 126, 389–402.
- Ward, K.M., Aletras, A.H., Balaban, R.S., 2000. A new class of contrast agents for MRI based on proton chemical exchange dependent saturation transfer (CEST). *J. Magn. Reson.* 143, 79–87.
- Watanabe, T., Radulovic, J., Spiess, J., Natt, O., Boretius, S., Frahm, J., Michaelis, T., 2001. In vivo 3D MRI staining of the mouse hippocampal system using intracerebral injection of MnCl₂. *NeuroImage* 22, 860–867.
- Weissleder, R., Simonova, M., Bogdanova, A., Bredow, S., Enochs, W.S., Bogdanov Jr., A., 1997. MR imaging and scintigraphy of gene expression through melanin induction. *Radiology* 204, 425–429.
- Yang, C.T., Li, Y., Liu, S., 2007. Synthesis and structural characterization of complexed of a DO3A-conjugated triphenylphosphonium cation with diagnostically important metal ions. *Inorg. Chem.* 46, 8988–8997.
- Yang, P.F., Chen, D.Y., Hu, J.W., Chen, J.H., Yen, C.T., 2011. Functional tracing of medial nociceptive pathways using activity-dependent manganese-enhanced MRI. *Pain* 152, 194–203.
- Yu, X., Wadghiri, Y.Z., Sanes, D.H., Turnbull, D.H., 2005. In vivo auditory mapping in mice with Mn-enhanced MRI. *Nat. Neurosci.* 8, 961–968.
- Yu, X., Sanes, D.H., Aristizabal, O., Wadghiri, Y.Z., Turnbull, D.H., 2007. Large scale reorganization of the tonotopic map in mouse auditory midbrain revealed by MRI. *Proc. Natl. Acad. Sci. USA* 104, 12193–12198.
- Zhang, S., Winter, P., Wu, K., Sherry, A.D., 2001. A novel europium(III)-based MRI contrast agent. *J. Am. Chem. Soc.* 123, 1517–1518.
- Zhang, X.A., Lovejoy, K.S., Jasanoff, A., Lippard, S.J., 2007. Water-soluble porphyrins as a dual-function molecular imaging platform for MRI and fluorescence zinc sensing. *Proc. Natl. Acad. Sci.* 104, 10780–10785.
- Zhang, X., Bearer, E.L., Boulat, B., Hall, F.S., Uhl, G.R., Jacobs, S.E., 2010. Altered neurocircuitry in the dopamine transporter knockout mouse brain. *PLoS One* 5, 11506.
- Ziv, K., Meir, G., Harmein, A., Shimoni, E., Klein, E., Neeman, M., 2010. Ferritin as a reporter gene for MRI: chronic liver over expression of H-ferritin during dietary iron supplementation and aging. *NMR Biomed.* 23, 523–531.
- Zurkiya, O., Chan, A.W., Hu, X., 2008. MagA is sufficient for producing magnetic nanoparticles in mammalian cells, making it an MRI reporter. *Magn. Reson. Med.* 59, 1225–1231.